



# “A Handbook of Integer Sequences” Fifty Years Later

N. J. A. Sloane

**B**efore 1973 there was no database of integer sequences. Someone coming across the sequence 1, 2, 4, 9, 21, 51, 127, ... would have had no way of discovering that it had been studied since 1870 (today these are called the Motzkin numbers, and they form entry [A001006](#) in the database). Everything changed in 1973 with the publication of *A Handbook of Integer Sequences*, which listed 2372 entries. This article describes the 50-year evolution of the database from the *Handbook* to its present form as the *On-Line Encyclopedia of Integer Sequences* (or *OEIS*), which contains 360,000 entries, receives a million visits a day, and has been cited 10,000 times, often with a comment saying “discovered thanks to the OEIS.”

## Integer Sequences

Number sequences arise in all branches of science: for example, 1, 1, 2, 4, 9, 20, 48, 115, ... gives the number of rooted trees with  $n$  nodes ([A000081](#); see also Figure 1). And for an example from daily life, into how many pieces can you cut a pancake with  $n$  knife-cuts? The pieces need not all be the same size. That one is easy: 1, 2, 4, 7, 11, 16, ...,  $n(n+1)/2 + 1$  ([A000124](#)). But what is the answer for cutting up an (ideal) bagel or doughnut? That is a lot harder: with a sharp knife you might get a few terms, perhaps 1, 2, 6, 13, ..., but probably not enough to guess the formula, which is  $n(n^2 + 3n + 8)/6$  for  $n > 0$ . For that you would need to consult the database: go to <https://oeis.org> and enter “cutting bagel,” or go directly to [A003600](#).

My fascination with these sequences began in 1964 when I was a graduate student at Cornell University, in Ithaca, New York, studying neural networks. I had encountered a sequence of numbers, 1, 8, 78, 944, 13800, ..., and I badly needed a formula for the  $n$ th term in order to determine the rate of growth of the sequence (this would indicate how long the activity in this very simple neural network would persist).

I noticed that although several books in the Cornell library contained sequences somewhat similar to mine, as far as I could tell, this particular sequence was not mentioned. I expected to have to analyze many related sequences, so

in order to keep track of the sequences in these books, I started recording them on 3" × 5" file cards.<sup>1</sup>

The collection grew rapidly as I searched through more books, and once the word got out, people started sending me sequences. Richard Guy was an enthusiastic supporter right from the start. In 1973, I formalized the collection as *A Handbook of Integer Sequences*, which was published by Academic Press (Figure 2).

Once the book appeared, the flood of correspondence increased, and it took 20 years to prepare the next version. Simon Plouffe helped a great deal, and in 1995, Academic Press published our sequel, *The Encyclopedia of Integer Sequences*, with 5487 entries. From this point on, the collection grew even more rapidly. I waited a year, until it had doubled in size, and then put it on the internet, calling it the *On-Line Encyclopedia of Integer Sequences*.

In the rest of this article I will first say more about the evolution of the database: the *Handbook*, the 1995 *Encyclopedia*, the *On-Line Encyclopedia*, and the *OEIS Foundation*. The next sections describe the database itself: what sequences are—or are not—included, how the database is used, the layout of a typical entry, the arrangement of the entries, and a fact sheet. The final sections describe some especially interesting sequences: Recamán’s sequence, the iteration of number-theoretic functions, Gijswijt’s sequence, lexicographically earliest sequences, the stepping stones problem, and stained glass windows. These last sections mention several open questions to which I would very much like to know the answers.

Notation:  $a(n)$  denotes the  $n$ th term of the sequence under discussion;  $\sigma(n)$  is the sum of the divisors of  $n$  ([A000203](#)).


## Evolution of the Database

### *The Handbook of Integer Sequences*


Once the collection had grown to a few hundred entries, I entered them on punched cards,<sup>2</sup> which made it easier to check and sort them. The *Handbook* was typeset directly from the punched cards. There were a few errors in the

<sup>1</sup>Six-digit numbers prefixed by A refer to entries in the current version of the *Handbook*, the *On-Line Encyclopedia of Integer Sequences* [12].

<sup>2</sup>These were never called “punch cards.” To anyone who worked with them in the 1960s, “punch cards” sounds like “grill cheese” for “grilled cheese” or “barb wire” for “barbed wire,” both of which I have recently seen in print.



**Figure 1.** (a) One of 48 unlabeled rooted trees with 7 nodes (the root node is at the bottom). (b) Four cuts of a pancake can produce 11 pieces. (c) Three cuts of a bagel can produce 13 pieces.



**Figure 2.** Front cover of the *Handbook*. The embossed figures show side views of the two ways of folding a strip of three (blank) stamps and the five ways of folding a strip of four stamps. The full sequence begins 1, 1, 2, 5, 14, 38, 120, 353, 1148, 3527, ..., [A001011](#). No formula is known.

book, but almost all of them were caused by errors in the original publications. Accuracy was a primary concern in that book, as it is today in the OEIS.

The book was an instant success. It was, I believe, the world's first dictionary of integer sequences (and my original title had *Dictionary* rather than *Handbook*). Many people said "What a great idea" and wondered why no one had done it before. Martin Gardner recommended it in the *Scientific American* of July 1974. Lynn A. Steen, writing in the *American Mathematical Monthly*, said:

Incomparable, eccentric, yet very useful. Contains thousands of "well-defined and interesting" infinite integer sequences together with references for each ... If you ever wondered what comes after 1, 2, 4, 8, 17, 35, 71, ..., this is the place to look it up.

Harvey J. Hindin, writing from New York City, exuberantly concluded a letter to me by saying, "There's the *Old Testament*, the *New Testament*, and the *Handbook of Integer Sequences*."

I never did find the sequence that started it all in the literature, but I learned Pólya's theory of counting, and with John Riordan's help found the answer, which appears in [14] and [A000435](#).

### *The Encyclopedia of Integer Sequences*

Following the publication of the *Handbook*, a large amount of correspondence ensued, with suggestions for further sequences and updates to the entries. By the early 1990s, over a cubic meter of new material had accumulated. A Canadian mathematician, Simon Plouffe, offered to help in preparing a revised edition of the book, and in 1995, *The Encyclopedia of Integer Sequences*, by me and Plouffe, was published by Academic Press. It contained 5487 sequences, occupying 587 pages. By now, punched cards were obsolete, and the entries were stored on magnetic tape.

### *The On-Line Encyclopedia of Integer Sequences*

Again, once the book appeared, many further sequences and updates were submitted from people all over the world. I waited a year, until the size of the collection had doubled, to 10,000 entries, and then in 1996, I launched the *On-Line Encyclopedia of Integer Sequences* (now usually called simply the *OEIS*) on the internet. From 1996 until October 26, 2009, it was part of my homepage on the AT&T Bell Labs website.

Incidentally, in 2004, the database was mentioned by the internet website *slashdot* ("News for Nerds. Stuff that Matters"), and this brought so much traffic to my AT&T Labs homepage that it briefly crashed the whole AT&T Labs website. My boss was quite proud of this, since it was a rare accomplishment for the Mathematics and Statistics Research Center.

### *The OEIS Foundation*

In 2009, in order to ensure the long-term future of the database, I set up a nonprofit foundation, the OEIS Foundation

Inc., a 501(c)(3) public charity, whose purpose is to own, maintain, and raise funds to support the *On-Line Encyclopedia of Integer Sequences*, or *OEIS*.

On October 26, 2009, I transferred the intellectual property of the *On-Line Encyclopedia of Integer Sequences* to the foundation. A new OEIS with multiple editors was launched on November 11, 2010.

Since then, it has been possible for anyone in the world to propose a new sequence or an update to an existing sequence. To do this, users must first register, and then submissions are reviewed by the editors before they become a permanent part of the OEIS. Technically, the OEIS is now a “moderated wiki.”

I started writing this article on November 11, 2022, noting that this marked 11 years of successful operation of the online OEIS, and that the database is in its 59th year of existence.

## The Database Today What Sequences Are Included?

From the very beginning, the goal of the database has been to include all “interesting” sequences of integers. This is a vague definition, but some further examples may make it clearer. The database includes a huge number of familiar and unfamiliar sequences from mathematics (the prime numbers, 2, 3, 5, 7, 11, 13, ..., [A000040](#); the orders of noncyclic simple groups, 60, 168, 360, 504, 660, 1092, ..., [A001034](#)); computer science (the number of comparisons needed for merge sort, 0, 1, 3, 5, 8, 11, 14, ..., [A001855](#)); physics (see “self-avoiding walks on lattices,” Ising model, etc., e.g., [A002921](#)); chemistry (the enumeration of chemical compounds was one of the motivations behind Pólya’s theory of counting; see, e.g., [A000602](#)); and not least, from puzzles and IQ tests: 1, 8, 11, 69, 99, 96, 111, ..., the “strobogrammatic” numbers, guess!, or see [A000787](#); 4, 14, 23, 34, 42, 50, 59, ..., the stops with numerical values for the A Train (8 Avenue express) in Manhattan, as of January 2023, [A011554](#). The latter entry has links to a map and the train schedule.

Sequences that have arisen in the course of someone’s work—especially if published—have always been welcomed. On the other hand, sequences that have been proposed simply because they were missing from the database are less likely to be accepted. And of course, the sequence must be well defined.

Very short sequences and sequences that are subsequences of many other sequences are not accepted. A sequence for which the only known terms are 2, 3, 5, 7 would not be accepted, since it is matched by a large number of existing sequences. The definition may not involve an arbitrary but large parameter (primes ending in 1 are fine, [A030430](#), but not primes ending in 2023).

The OEIS [Wiki](#) has a section listing additional examples of what not to submit, as well as a great deal of information about the database that I won’t repeat here, such as the meaning of the various keywords, the definition of the “offset” of a sequence, descriptions of the submission and editorial processes, and a list of over 10,000 citations of the OEIS in the scientific literature.

Most OEIS entries give an ordered list of integers. But triangles of numbers are included by reading them row by row. For example, Pascal’s triangle becomes 1, 1, 1, 1, 2, 1, 1, 3, 3, 1, ..., [A007318](#). Doubly infinite square arrays are included by reading them by antidiagonals: the standard multiplication table for positive integers becomes 1, 2, 2, 3, 4, 3, 4, 6, 6, 4, ..., [A003991](#).

Sequences of fractions are included as a linked pair giving the numerators and denominators separately (the Bernoulli numbers are [A027641/A027642](#)). Important individual real numbers are included by giving their decimal or continued fraction expansions (for  $\pi$ , see [A000796](#) and [A001203](#)). A relatively small number of sequences of nonintegral real numbers are included by rounding them to the nearest integer or by taking floors or ceilings (the imaginary parts of the nontrivial zeros of Riemann’s zeta function give [A002410](#)).

Two less obvious sources for sequences are binomial coefficient identities and number-theoretic inequalities. The values of either side of the identity

$$\sum_{k=0}^n \binom{2n}{k}^2 = \frac{1}{2} \binom{4n}{2n} - \frac{1}{2} \binom{2n}{n}^2$$

[8, (3.68)] give [A036910](#). From the inequality  $\sigma(n) < n\sqrt{n}$  for  $n > 2$  [11, Sect. III.1.1.b], we get the integer sequence  $\lfloor n\sqrt{n} \rfloor - \sigma(n)$ , [A055682](#). The point here is that if you want to find out whether this inequality is known, you look up the difference sequence, and find [A055682](#) and a reference to the proof. Many more sequences of these two types should be added to the database.

## How the Database Is Used

The main applications of the database are in identifying sequences and in finding out the current status of a known sequence. Barry Cipra has called it a mathematical analogue of a “fingerprint file.” You encounter a number sequence and wish to know whether anyone has ever come across it before. You enter the first few terms of the sequence into the search field on the OEIS home page, and if your sequence is in the database, the reply will give a definition of the first 50 or so terms, and, when available, formulas, references, computer code for producing the sequence, links to any relevant web sites, and so on.

Figures 3 and 4 show what happens if you enter 1, 2, 5, 14, 42, 132, 429, the first few Catalan numbers, one of the most famous sequences of all.

I could have chosen a simpler example, like the Fibonacci numbers, but I have a particular reason for choosing the Catalan numbers. When the OEIS was new, people would sometimes say to me that they had a sequence they were trying to understand, and would I show them how to use the database. At least twice when I used the Catalan sequence as an illustration, they said, “Why, that is my sequence! How on earth did you know?” It was no mind-reading trick. The Catalan numbers are certainly the most common sequence that people don’t know about. This entry is the longest—and one of the most important—in the whole database.

The OEIS is supported by [the many generous donors to the OEIS Foundation](#).

0 1 3 6 2 7  
 : 13  
 : 20  
 23  
 10 22 11 21

THE ON-LINE ENCYCLOPEDIA  
 OF INTEGER SEQUENCES®

founded in 1964 by N. J. A. Sloane

1, 2, 5, 14, 42, 132, 429  [Hints](#)

(Greetings from [The On-Line Encyclopedia of Integer Sequences!](#))

Search: **seq:1,2,5,14,42,132,429**

Displaying 1-10 of 26 results found. page 1 [2](#) [3](#)

Sort: relevance | [references](#) | [number](#) | [modified](#) | [created](#)    Format: long | [short](#) | [data](#)

**Figure 3.** The result of querying the database with 1, 2, 5, 14, 42, 132, 429. This figure shows the banner at the top of the reply. There are 26 matches, ranked in order of importance, the top match being the one we want, the Catalan numbers. A shortened version of the top match is shown in the next figure.

A000108	Catalan numbers: $C(n) = \text{binomial}(2n, n)/(n+1) = (2n)!/(n!(n+1)!)$ . (Formerly M1459 N0577)
DATA	1, 1, 2, 5, 14, 42, 132, 429, 1430, 4862, 16796, 58786, ...
COMMENTS	These were formerly sometimes called Segner numbers. A very large number of combinatorial interpretations are known - see references, esp. R. P. Stanley, Catalan Numbers, Camb., 2015. This is probably the longest entry in the OEIS, and rightly so. The solution to Schröder's first problem: number of ways to insert $n$ pairs of parentheses in a word of $n+1$ letters. E.g., for $n=2$ there are 2 ways: ((ab)c) or (a(bc)); for $n=3$ there are 5 ways: ((ab)(cd)), (((ab)c)d), ...
REFERENCES	... The large number of references and links demonstrates the ubiquity of the Catalan numbers. R. Alter, Some remarks and results on Catalan numbers, pp. 109-132 in Proc. Louisiana Conf. Combinatorics, Graph Theory and Computer Science. Vol. 2, edited R. C. Mullin et al., 1971. M. Bona, ed., Handbook of Enumerative Combinatorics, CRC Press, 2015 L. Comtet, Advanced Combinatorics, Reidel, 1974, p. 53. J. H. Conway & R. K. Guy, The Book of Numbers, Springer, 1995, 96-106. ...
LINKS	Robert G. Wilson v, Table of $n, a(n)$ for $n = 0..1000$ ... F. R. Bernhart, Catalan, Motzkin and Riordan numbers, Discr. Math., Vol. 204, No. 1-3 (1999), 73-112. ... W. G. Brown, Historical Note on a Recurrent Combinatorial Problem, Amer. Math. Monthly, 72, No. 9 (1965), 973-977. ...
FORMULA	Recurrence: $a(n) = 2 * (2 * n - 1) * a(n - 1)/(n + 1)$ with $a(0) = 1$ . ...
MAPLE	A000108 := n->binomial(2*n, n)/(n+1); ...
MATHEMATICA	A000108[n_] := (2 n)!/n!/(n+1)!
PARI	a(n)=binomial(2*n, n)/(n+1) ...
KEYWORD	core,nonn,easy,nice
AUTHOR	N. J. A. Sloane

**Figure 4.** The entry for the Catalan numbers [A000108](#). The full entry has over 750 lines, which have been edited here to show samples of the different fields.

If we do not find your sequence in the database, we will send you a message inviting you to submit it (if you consider it to be of general interest), so that the next person who comes across it will be helped, and your name will go on record as the person who submitted it.

The second main use of the database is to find out the latest information about a particular sequence. Of course, we cannot hope to keep all 360,000 entries up-to-date. But when a new paper is published that mentions the OEIS, Google will tell us, and we then add links to that paper from every sequence that it mentions. People have told us that this is one of the main ways they use the OEIS. After all, even a specialist in (say) permutation groups cannot keep track of all the papers published worldwide in that area. And if a paper in a physics or engineering journal happens to mention a number-theoretic sequence, for example, Google will notify us and we will record it.

There are also many other ways in which the database has proved useful. For example, it is an excellent source of problems to work on. The database is constantly being updated. Every day, we get thirty to fifty submissions of new sequences, and an equal number of comments on existing entries (new formulas, references, additional terms, etc.). The new sequences are often sent in by nonmathematicians, and they are a great source of problems. You can see the current submissions at <https://oeis.org/draft>. Often enough, you will see a sequence that is so interesting you want to drop everything and work on it. And remember that we are always in need of more volunteer editors. In fact, anyone who has registered with the OEIS can suggest edits; you do not even need to be an official editor. We have been the source of many international collaborations.

There is also an educational side: several people have told us that they were led into mathematics through working as an editor. Here is a typical story.

Subject: Reminiscence from a young mathematician

I wanted to relay a bit of nostalgia and my heartfelt thanks. Back in the late 1990s, I was a high school student in Oregon. While I was interested in mathematics, I had no significant mathematically creative outlet until I discovered the OEIS in the course of trying to invent some puzzles for myself. I remember becoming a quite active contributor through the early 2000s, and eventually at one point, an editor. My experience with the OEIS, and the eventual intervention of one of my high school teachers, catalyzed my interest in studying mathematics, which I eventually did at ... College. I went on to a Ph.D. in algebraic geometry at the University of ... and am currently at ....

I wanted to thank you for seriously engaging with an 18-year-old kid, even though I likely submitted my fair share of mathematically immature sequences. I doubt I would have become a mathematician without the OEIS!

A less obvious use of the database is to quickly tell you how hard a problem is. I use it myself in this way all the time. Is the sequence “Catalan” or “Collatz”? If a sequence

comes up in your own work or when you are reviewing someone else’s work, it is useful to know right away whether it is a well-understood sequence, like the Catalan numbers, or whether it is one of the notoriously intractable problems like the Collatz, or  $3x + 1$ , problem ([A006577](#)).

Finally, the OEIS is a welcome escape when you feel the world is falling apart. Take a look at Scott Shannon’s drawings of stained glass windows in [A331452](#) or Jonathan Wild’s delicate illustrations of the ways to draw four circles in [A250001](#) or Éric Angelini’s “1995” puzzle ([A131744](#)) or any of his “lexicographically earliest sequences” ([A121053](#), [A307720](#), and many more); or find better solutions to the Stepping Stones Problem ([A337663](#)). You can find brand new problems at any hour of the day or night by looking at the stack of recent submissions: but beware, you may see a problem there that will keep you awake for days. Or search in the database for phrases like “It appears that ...” or “Conjecture: ...” or “It would be nice to know more!”

### Layout of a Typical Entry

This is a good place to mention some of the features of an OEIS entry. Most of the fields (see Figures 3 and 4) are self-explanatory. At the top, it tells you how many matches were found to your query (26 in the example). These are ranked in order of importance.

The DATA section shows the start of the sequence, usually enough terms to fill a few lines on the screen (typically 300 to 500 decimal digits). All terms listed must be known to be correct, and there cannot be any gaps. If the first  $n$  terms are known but the  $(n + 1)$ st is known only to be either 14 or 15, then the listing of the sequence must end with the  $n$ th term. In the case of Mersenne primes ([A000043](#)), it is common for later primes to be discovered before all smaller candidates have been tested. Until it is known for certain that the new discoveries are indeed the *next* terms, they cannot be added to the sequence (although of course they can be mentioned in comments). Often, one wants more terms than are shown in the DATA section, and in many cases, the first link in the entry will point to a plain-text file with perhaps 10,000 or 20,000 terms. That file will have a name like `b001006.txt` and is called the “b-file” for the sequence. Some entries also have much larger tables, giving a million or more terms.

If you click the “graph” button near the top of the reply, you will be shown two plots of the sequence, and if you click the “listen” button, you can listen to the sequence played on an instrument of your choice. The default instrument is the grand piano, and the terms of the sequence would then be mapped to the 88 keys by reducing the numbers mod 88 and adding 1.

I conclude this section with a philosophical comment. When you are seriously trying to analyze a sequence and are prepared to spend any amount of time needed (searching for a formula or recurrence, for instance), you need all the help you can get, which is why we provide the b-files and other data files, and why we give computer programs in so many languages. This is also the reason we give as many references and links as possible for a sequence. Even if the reference is to an ancient or obscure journal, or one

that has been accused of being “predatory,” we still give the reference, especially for sequences that are not well understood. The same thing holds for formulas, comments, and cross references to other sequences. When you are desperate, you will accept help from anywhere. And do not forget “[Superseeker](#),” which invokes a collection of algorithms to try to analyze your sequence or to transform it into an OEIS entry.

## Arrangement of the Entries

The entries in the database are (virtually) arranged in two different ways, the first essentially chronological, the second lexicographic.

The first is by an entry’s *absolute* identification number, or A-number.<sup>3</sup> Once the collection reached a few hundred entries, I sorted them into lexicographic order and numbered them A1, A2, A3, .... The sequence A1 gives the number of symmetry groups of order  $n$ , A2 is the famous Kolakoski sequence, and so on. This numbering is still used today, only A1 has become [A000001](#), A2 is [A000002](#), and as each new submission comes in, it gets the next free number. Currently, sequences are being issued numbers around A360000. Rejected A-numbers are recycled, so there are no gaps in the order. We reached 100,000 entries in 2004, and 250,000 in 2015. The present growth rate is about 12,000 new entries each year.

The second arrangement is a kind of lexicographic ordering. First I describe an idealized, theoretical, lexicographic order. Sequences of nonnegative numbers can be arranged in lexicographic (or dictionary) order. For example, sequences beginning 1, 2, 4, ... come before 1, 2, 5, ..., 1, 3, ..., etc., but after 1, 2, 3, .... Also, 1, 2, 4, ... comes after the two-term sequence 1, 2, because blanks precede numbers.

More formally, we compare the two sequences term by term, and in the first position where they differ, whichever is smaller (or blank) is the lexicographically earlier one. For sequences with negative terms, we ignore the signs and sort according to the absolute values.

Here is the actual ordering used in the OEIS. The sequences are arranged (virtually) into a version of lexicographic order, according to the following rules. First, delete all minus signs. Then find the first term that is greater than 1 and discard all the terms before it. What’s left determines its position in the lexicographic order. For example, to place  $-1, 0, 1, 1, \underline{2}, 1, 17, -3, -2, 6, \dots$  in the ordering, we would ignore the terms before the underlined 2 and consider the sequence as beginning 2, 1, 17, 3, 2, 6, ....

Sequences that contain only 0’s, 1’s and  $-1$ ’s are sorted into lexicographic order by absolute value and appear at the beginning of the ordering. The first sequence in the database is therefore the zero sequence [A000004](#).

In this way, every sequence has a unique position in the ordering. The sequences have been sorted in this way since the 1960s. For the first 10 years, the punched card entries were physically sorted into this order.

When you look at any OEIS entry, [A005132](#) say, toward the bottom you will see<sup>4</sup> two lines like

```
Sequence in context: A277558 A350578 A335299
                    * A064388 A064387 A064389
Adjacent sequences: A005129 A005130 A005131
                    * A005133 A005134 A005135
```

which tell you the three entries immediately before and after that entry in the lexicographic ordering and the three entries before and after it in the A-numbering. The asterisks represent the sequence you are looking at. The first group can be useful if you are uncertain about a term in your sequence, the second in case you want to look at other sequences submitted around that time.

Today, the sequences are stored internally in an SQLite database. However, the punched card format has been so useful that when you view a sequence, as in Figure 4, it is still presented to you in something very like the old punched card format.

## Summary: “A Handbook of Integer Sequences” Today

- Now the *On-Line Encyclopedia of Integer Sequences* or *OEIS*: <https://oeis.org>
- Accurate information about 360,000 sequences.
- Definition, formulas, references, links, programs. View as list, table, graph, music!
- Traffic: 1 million hits a day.
- 30 new entries, 50 updates every day.
- Often called one of best math sites on the Web. Fingerprint file for mathematics.
- “Street creds”: [10,000 citations](#).
- A moderated Wiki, owned by OEIS Foundation, a 501(c)(3) public charity.
- Uses: to see whether your sequence is new, to find references, formulas, programs.
- Catalan or Collatz? (Very easy or very hard?)
- Source of fascinating research problems;<sup>5</sup> low-hanging fruit from recent submissions.
- Accessible (free, friendly).
- Fun (1, 2, 4, 6, 3, 9, 12, 8, 10, 5, 15, ...?). Interesting, educational. Escape.
- Addictive (better than video games).
- Has led many people into mathematics.
- One of the most successful international collaborations, a modest contribution toward world peace.
- Needs editors.


## Some Favorite Sequences

I am sometimes asked for my favorite sequence. This is a difficult question. I’m tempted to reply by saying, “If you were the keeper of the only zoo in the world, how would

<sup>3</sup>The sequences in the 1973 and 1995 books were numbered respectively N0001, ... and M0001, ....

<sup>4</sup>If you don’t see them, click on the A-number at the top of the entry.

<sup>5</sup>Look for “Conjecture,” “It appears that,” “It would be nice to,” ...



**Figure 5.** Harriss's drawing of the first 64 terms of Recamán's sequence. (The tiny initial semicircle, at the extreme left, is below the axis. It has diameter 1 and joins the points 0 and 1. It continues as a semicircle of diameter 2 above the axis, joining the points 1 and 3.)

you answer that question?" (Because that is roughly the situation I'm in.) Would you pick one of the exotic animals, a giraffe, a kangaroo, or a blue whale? Or one of the essential animals, like a horse, a cow, or a duck? If the question came from a visiting alien, then of course there would be only one possible answer: a human being.

For sequences, the essential ones are the primes, the powers of 2, the Catalan numbers, and (especially if the question came from an alien with no fingers or toes) the counting sequence 0, 1, 2, 3, 4, ... ([A001477](#)).

But here I'll mention a few that are fairly exotic. The Recamán and Gijswijt sequences have simple recursive definitions, yet are astonishingly hard to understand.

### Recamán's Sequence ([A005132](#))

This remarkable sequence has resisted analysis for over 30 years, even though we have computed an astronomical number of terms. It was contributed to the database by Bernardo Recamán Santos in 1991.

The definition is deceptively simple. The first term is 0. We now add or subtract 1, then we add or subtract 2, then add or subtract 3, and so on. The rule is that we always first try to subtract, but we can subtract only if that leaves a nonnegative number that is not yet in the sequence. Otherwise, we must add.

Here is how the sequence begins. We have the initial 0. We can't subtract 1, because that would give a negative number, so we add 1 to 0. So the second term is 1. We can't subtract 2 from 1, so we add it, getting the third term  $1 + 2 = 3$ . Again we can't subtract 3, for that would give 0, which has already appeared, so we add 3, getting the fourth term  $3 + 3 = 6$ .

Now we must add or subtract 4, and this time we can subtract, because  $6 - 4 = 2$ , and 2 is nonnegative and a number that hasn't yet appeared. So at this point, the sequence is 0, 1, 3, 6, 2. Then it continues with  $7 (= 2 + 5)$ ,  $13 (= 7 + 6)$ ,  $20 (= 13 + 7)$ ,  $12 (= 20 - 8)$ , and so on. The terms  $a(0)$  through  $a(15)$  are

0, 1, 3, 6, 2, 7, 13, 20, 12, 21, 11, 22, 10, 23, 9, 24.

When you are adding rather than subtracting, repeated terms are permitted (e.g.,  $a(20) = a(24) = 42$ ).

Edmund Harriss has found an elegant way to draw the sequence as a spiral on the number line (Figure 5). Start at 0, and when we subtract  $n$ , draw a semicircle of diameter  $n$  to the left from the last point, and draw one to the right when we are adding  $n$ . Draw the semicircles alternately below and above the horizontal axis so as to produce a smooth spiral.

The main question about this sequence is this: does every positive number appear? What makes this sequence so interesting is that certain numbers (for reasons we do not understand) are extremely reluctant to appear. For example, 4 does not appear until 131 steps, and 19 takes 99,734 steps.

A group of us at AT&T Labs worked on this in 2001 and found a way to greatly speed up the computation. Allan Wilks used our method to compute the first  $10^{15}$  terms and found that 2406 (which had been missing for a long time) finally appeared at step 394,178,473,633,984.

At that point, the smallest missing number was  $852655 = 5 \cdot 31 \cdot 5501$ . Benjamin Chaffin has continued this work, and in 2018, he reached  $10^{230}$  terms. However, 852,655 was still missing, and there has been no progress since then.

Thirty years ago, I thought that every number would eventually appear. Now I am not so sure. My current belief is that there are two possibilities: (1) There are infinitely many numbers that never appear, and 852,655 just happens to be the smallest of them and has no other special property. A similar phenomenon seems to occur in iterating various number-theoretic functions—see the next section. (2) Every number will eventually appear (just as presumably every one of Shakespeare's plays will eventually appear in the expansion of  $\pi$  in base 60), although we may never be able to extend the sequence far enough to hit 852,655. For the latest information about this sequence (or any other sequence mentioned in this article), consult the OEIS.

Open question: Does 852,655 appear in [A005132](#)?

### Iteration of Number-Theoretic Functions

Many mysterious sequences arise from the iteration of number-theoretic functions. A classic problem concerns the iteration of the function  $f(n) = \sigma(n) - n$ , the sum of the divisors of  $n$  that are less than  $n$  (technically, the sum of the "aliquot parts" of  $n$ ) [A001065](#). For an initial value of  $n$ , what happens to the sequence (or "trajectory")  $n, f(n), f(f(n)), f(f(f(n))), \dots$ ? All  $n < 276$  terminate by entering a cycle (such  $n$  are called "perfect," "amicable," or "sociable" numbers, depending on whether the cycle is of length 1, 2, or  $\geq 3$ ) or by reaching a prime, then 1, then 0.

But it appears likely that  $n = 276$ , and perhaps all sufficiently large even numbers, will never terminate [[5](#), [9](#), §B6]. The trajectory of 276 is sequence [A008892](#). At the time of writing, 2145 terms of this trajectory have been computed, and it is still steadily growing, term 2145 being a 214-digit number [[7](#)].

There are arguments that suggest that 276 will eventually terminate, and other arguments that suggest it will grow forever. It is surprising that even today, mathematics cannot resolve such a concrete question.

Sequence [A098007](#) gives the number of distinct terms in the trajectory of a general initial number  $n$ , or  $-1$  if the trajectory is unbounded. The value of [A098007\(276\)](#) is unknown.

If indeed 276 does go to infinity, it is natural to ask, how did 276 know it was destined to be the first immortal number under the map  $f$ ? The answer may be that there are infinitely many immortal numbers, and 276 just happens to be the first. It got lucky, that's all! Just as 852,655 got lucky in Recamán's problem.

A similar question, discussed by Richard K. Guy [9, §B41], which has received much less attention, concerns the map  $g(n) = (\sigma(n) + \phi(n))/2$ , where  $\phi(n)$  is the Euler totient function [A000010](#). The trajectory may end at 1, a prime, or a fraction, or it may increase monotonically to infinity. Sequence [A292108](#) gives the number of steps in the trajectory, or  $-1$  if the trajectory is infinite. All numbers  $n < 270$  have finite trajectories, but it appears that 270 increases forever. The trajectory of 270 is [A291789](#). For this problem there is less doubt about what happens, because Andrew Booker has given a heuristic argument showing that almost all numbers go to infinity. What makes 270 the first immortal number under  $g$ ? Again I suspect it just got lucky!

Open questions: Does the trajectory of 276 under  $f$  increase forever? What about the trajectory of 270 under  $g$ ?

### Gijswijt's Sequence ([A090822](#))

For this sequence, it will be helpful to remember that chemists do not write HHO for water; they write  $H_2O$ . And they do not write AlAlAlS0000S0000; they write  $Al_3(SO_4)_2$ . We will apply a similar compression to sequences of numbers, except that we indicate repetition by superscripts rather than subscripts.

For this problem, when we look at a sequence of numbers, we want to write it in the form  $XY^k$ , where  $X$  and  $Y$  are themselves sequences of numbers,  $X$  can be missing, and the exponent  $k$  is *as large as possible*.

For example, we can write 1, 2, 2, 2, 2 as  $XY^k$ , where  $X = 1$ ,  $Y = 2$ , and  $k = 4$ . The highest  $k$  we can achieve for a sequence is called its curling number. So 1, 2, 2, 2, 2 has curling number 4. If you think of an animal with its head looking to the left, with a very curly tail, then  $X$  represents the head and body of the animal, and  $Y^k$  represents the curls in its tail.

Consider the sequence 3, 2, 4, 4, 2, 4, 4, 2, 4, 4. We could take  $X = 3, 2, 4, 4, 2, 4, 4, 2$  and  $Y = 4$ , getting  $XY^2$ , with  $k = 2$ , or we could take  $X = 3, Y = 2, 4, 4$ , getting  $XY^3$ , with  $k = 3$ , which is larger. So this sequence has curling number 3.

Remember that  $X$  may be missing. So the sequence with a single term 99, say, can be written as  $Y^1$ , where  $Y$  is the number 99, and it has curling number 1. The notion of curling number is independent of the base in which the numbers are written.

We are now ready to define Dion Gijswijt's absolutely brilliant sequence, which he sent to the OEIS in 2004.

The rule for finding the next term is simple: it is the curling number of the sequence so far. And you start with 1. That's the sequence!

So let's construct it. We start with 1, and the curling number of 1 is 1. So now we have 1, 1. This has curling number 2, so now we have 1, 1, 2. At each step we recompute the curling number and make that the next term.

Here are the first few generations:

```

1
1 1
1 1 2
1 1 2 1
1 1 2 1 1
1 1 2 1 1 2
1 1 2 1 1 2 2
1 1 2 1 1 2 2 2
1 1 2 1 1 2 2 2 3

```

To go from line 6 to line 7, we took  $Y = 1, 1, 2$ . In line 8, we see the first 3, at the ninth term, and after a while, a 4 appears at term 220.

But Gijswijt was unable to find a 5, and he left that question open when he submitted the sequence. Some AT&T Labs colleagues computed many millions of terms, but no 5 appeared.

Finally, over the course of a long weekend, Fokko van der Bult (a fellow student of Gijswijt's in Amsterdam) and I independently showed that there is a 5. In fact, there are infinitely many 5's, but the first one does not appear until about term  $10^{10^{23}}$ . The universe would be cold long before any computer search would find it.

In the paper we wrote about the sequence [15], we also conjectured that the first time a number  $N > 4$  appears is at about term

$$2 \uparrow (2 \uparrow (3 \uparrow (4 \uparrow (5 \uparrow \dots \uparrow (N - 1))))),$$

where the up-arrows ( $\uparrow$ ) indicate exponentiation, a tower of exponents of height  $N - 1$ . This is a very slow-growing sequence.

A very recent manuscript by a student of Gijswijt's, Levi van de Pol [16], still under review, has extended our work and may have proved the above conjecture.

I cannot resist adding a further comment about curling numbers, which if true shows that the Gijswijt sequence is in a sense universal. My "curling number conjecture" asserts that if any finite starting sequence is extended by the rule that the next term is the curling number of the sequence so far, then eventually the curling number will be 1.

If true, this implies that if the starting sequence contains no 1's, then the sequence eventually becomes Gijswijt's sequence [4, Theorem 23]. In fact, I conjecture that this is true for any starting sequence.

Open question: Is the curling number conjecture true?

## Lexicographically Earliest Sequences

Although there is no space here to discuss them in detail, there are many fascinating and difficult sequences in the OEIS whose definition has the form “Lexicographically Earliest Sequence of distinct positive numbers with the property that ...,” where now we are using lexicographic in its pure sense, as mentioned when we were discussing the arrangement of sequences in the database. A favorite example is the EKG (or ECG) sequence [A064413](#), whose definition is the lexicographically earlier infinite sequence of distinct positive numbers with the property that each term after the first has a nontrivial common factor with the previous term [10]. Other L.E.S. examples are the Yellowstone permutation [A098550](#) [2], the Enots Wolley sequence [A336957](#) (the name suggests the definition), and the Binary Two-Up sequence [A354169](#) [6].

Open question: show that the terms of the Enots Wolley sequence are precisely 1, 2, and all numbers with at least two distinct prime factors.

## The Stepping Stones Problem (A337663)

This lovely problem was invented in 2020 by two undergraduates, Thomas Ladouceur and Jeremy Rebenstock. You have an infinite chessboard and a handful of brown stones, which are worth one point each. You also have an infinite number of white stones, of values 2, 3, 4, ..., one of each value. Suppose you have  $n$  brown stones. You start by placing them anywhere on the board. Now you place the white stones, trying to place as many as you can. The rules are that you can place a white stone with value  $k$  on a square only if the values of the stones on the eight squares around it add up to  $k$ . And you must place the white stones in order, first 2, then 3, and so on. You stop when you cannot place the next-higher-numbered white stone. The goal is to find the highest value that can be placed. Call this value, for the game with  $n$  stones,  $a(n)$ .

Say we start with  $n = 2$  brown stones. There are infinitely many squares on which they can be placed, but in order for us to be able to place the white stone with value 2, they must be placed with at most one blank square between them. It turns out that the best thing is to place them so they are separated diagonally by a single blank square, as in Figure 6.

Now we start trying to place the white stones. The 2 stone has to go between the two brown (or 1) stones, and then the 3 goes on a square adjacent to the 1 and the 2. There is now a choice for where the 4 goes, but the choice shown in Figure 6 is the best. (After we have placed the 4, the neighbors of the 3 no longer add up to 3, but that is OK. It is only *when we place* the 3 that its neighbors must sum to 3.) Continuing in this way, we eventually reach 16. There is nowhere to place the 17, so we stop. Ladouceur and Rebenstock showed, using a computer and considering all possible arrangements, that 16 is the highest value that can be attained with two starting stones. So  $a(2) = 16$ .

	9	5	10	11			
		4	1				
	12	8	3	2		16*	
				6	1	15	
			13	7	14		

Figure 6. A solution to the Stepping Stones Problem for two starting stones. The high point  $a(2) = 16$  here is indicated by an asterisk, as it is in the next three tables.

	35	18	36		23		21		32					
		17	1		14	9		12	20					
		34	16	15		5	4	8			26	27		
				31		10	1	3	19	25		1	28	
						11		2	6		33			29
						24	13	22	1	7				
						37			30	38*				

Figure 7. A solution to the Stepping Stones Problem for four starting stones.

This is clearly a hard problem, since the number of possibilities grows rapidly with the number of brown stones. Only six terms of this sequence are known:  $a(1)$  through  $a(6)$  are 1, 16, 28, 38, 49, 60. A solution for  $n = 4$  found by Arnaud Chevallier is shown in Figure 7. There are lower bounds for larger values of  $n$  that may turn out to be optimal. For  $n = 7, 8, 9, 10$ , the current best constructions give 71, 80, 90, 99. See [A337663](#) for the latest information.

We don’t know how fast  $a(n)$  grows. Some upper and lower bounds, initiated by Robert Gerbicz and Andrew Howroyd, can be found in the Comments section of [A337663](#). The simple linear construction shown in Figure 8 shows that  $a(n) \geq 6(n - 1)$  for  $n \geq 3$ .

By combining the constructions of Figures 6 and 8, Menno Verhoeven obtained  $a(n) \geq 6n + 3$  for  $n \geq 3$  (Figure 9).

The best lower bound for large  $n$  is due to Robert Gerbicz, who has shown by a remarkable extension of the construction in Figures 8 and 9 that  $\lim_{n \rightarrow \infty} a(n)/n > 6$ .

(A preliminary version of his bound gives

$$a(n) > 6.0128n - 5621$$

for all  $n$ , although the exact values of the constants have not been confirmed.) In his construction, the “chimney” on the right of Figure 9 gets expanded into a whole trellis.

One might think that with a sufficiently clever arrangement, perhaps extending the construction in Figure 8 so that the path wraps around itself in a spiral, one could






Figure 11. A  $6 \times 6$  grid with every pair of boundary points joined by a chord. There are 4825 vertices and 6264 cells.

and [A255011](#) give the numbers of vertices and cells for  $n \leq 42$ . Sequence [A334699](#) enumerates the cells by number of sides.

In the summer of 2022, Scott Shannon and I considered several other families of planar graphs. I cannot resist showing one of Shannon's graphs, a  $16 \times 16$  grid, illustrating the 16th term of [A355798](#) (Figure 12). There are 61,408 cells. Although Shannon has calculated 40 terms of this sequence, again no formula is known.

### Other Sequences I Would Have Liked to Include

It is getting late, so I had better stop. Another time, I'll tell you about some very interesting sequences, such as those arising from the problem of dissecting a square to get a

regular  $n$ -gon ([A110312](#)); from gerrymandering ([A341578](#), [A348453](#), and many others); counting the ways in which circles can overlap ([A250001](#)); the inventory sequence ([A342585](#)); Kaprekar's junction numbers ([A006064](#), [1]); the kissing number problem ([A001116](#), [A257479](#)); the neural network problem that started it all ([A000435](#)); and the squares in the plane problem ([A051602](#)). And perhaps also metasequences such as [A051070](#) ( $a(n)$  is the  $n$ th term of  $A_n$ ) and [A107357](#) (the  $n$ th term is 1+ the  $n$ th term of  $A_n$ ).

A final comment: there are many videos in the internet of talks I have given about sequences, including several that Brady Haran has made for the Youtube Numberphile channel (which have been viewed over eight million times). See, for example, "[Terrific Toothpick Patterns.](#)"




Figure 12. Scott Shannon's "Magic Carpet" graph, illustrating [A355798\(16\)](#).

### Acknowledgments

I would like to thank some good friends who have helped me and the OEIS over the years: David L. Applegate, William Cheswick, Russ Cox, Susanna S. Cuyler, Harvey P. Dale, Ronald L. Graham, Richard K. Guy, Marc LeBrun, John Riordan, and Doron Zeilberger. There are many active volunteer editors, and it is impossible to thank them all. But I would like to give particular thanks to Jörg Arndt, Michael S. Branicky, Michael De Vlieger, Amiram Eldar, Charles R. Greathouse IV, Maximilian F. Hasler, Alois P. Heinz, Andrew Howroyd, Sean A. Irvine, Antti Karttunen, Michel Marcus, Richard J. Mathar, Peter Munn, Hugo Pfoertner, Kevin Ryde, Jon E. Schoenfeld, Rémy Sigrist, and Chai Wah Wu. I also thank the members of the Board of Trustees of the OEIS Foundation, past and present, for all their help, both to me personally and to the OEIS. Figure credits: Figure 1c: Clifford A. Pickover. Figure 5: Edmund Harriss. Figures 6, 7, 8, and 9 are based on communications from Thomas Ladouceur and Jeremy Rebenstock, Arnauld Chevallier, Skylark Xentha Murphy-Davies, and Menno Verhoeven, respectively. Figures 10 and 11: Lars Blomberg and Scott R. Shannon. Figure 12: Scott R. Shannon. Other figures: the author.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

### References

- [1] M. A. Alekseyev and N. J. A. Sloane. On Kaprekar's junction numbers. *J. Combinat. and Number Theory* 12:3 (2020) [2023], 115–155.

- [2] D. L. Applegate, H. Havermann, B. Selcoe, V. Shevelev, N. J. A. Sloane, and R. Zumkeller. The Yellowstone permutation. *J. Integer Seqs.* 18 (2015), #15.6.7.
- [3] L. Blomberg, S. R. Shannon, and N. J. A. Sloane. Graphical enumeration and stained glass windows, 1: rectangular grids. *Integers*, Ron Graham Memorial Volume 21A (2021), #A5.
- [4] B. Chaffin, J. P. Linderman, N. J. A. Sloane, and A. R. Wilks. On curling numbers of integer sequences. *J. Integer Seqs.* 16 (2013), #13.4.3.
- [5] K. Chum, R. K. Guy, M. J. Jacobson, Jr., and A. S. Mosunov. Numerical and statistical analysis of aliquot sequences. *Exper. Math.* 29:4 (2020), 414–425.
- [6] M. De Vlieger, T. Scheuerle, R. Sigrist, N. J. A. Sloane, and W. Trump. *The binary Two-Up sequence*. arXiv:2209.04108, 2022.
- [7] FactorDB database. Available at <http://factordb.com/sequences.php>, 2023.
- [8] H. W. Gould. *Combinatorial Identities*. Privately printed, Morgantown, West Virginia, 1972.
- [9] R. K. Guy. *Unsolved Problems in Number Theory*, 3rd ed. Springer, 2010.
- [10] J. C. Lagarias, E. M. Rains, and N. J. A. Sloane. The EKG sequence. *Exper. Math.* 11 (2002), 437–446.
- [11] D. S. Mitrinović, J. Sándor, and B. Crstici. *Handbook of Number Theory*. Kluwer, 1996.
- [12] OEIS Foundation Inc. *On-Line Encyclopedia of Integer Sequences*. Available at <https://oeis.org>, 2023.
- [14] B. Poonen and M. Rubinstein. The number of intersection points made by the diagonals of a regular polygon. *SIAM J. Discr. Math.* 11:1 (1998), 135–156.
- [15] J. Riordan and N. J. A. Sloane. Enumeration of rooted trees by total height. *J. Austral. Math. Soc.* 10 (1969), 278–282.
- [13] F. J. van de Bult, D. C. Gijswijt, J. P. Linderman, N. J. A. Sloane, and Allan Wilks. A slow-growing sequence defined by an unusual recurrence. *J. Integer Seqs.* 10 (2007), #07.1.2.
- [16] L. van de Pol. The first occurrence of a number in Gijswijt’s sequence. arXiv:2209.04657, 2022.

**N. J. A. Sloane**, The OEIS Foundation Inc., 11 S. Adelaide Ave., Highland Park, NJ 08904, USA.

E-mail: [njasloane@gmail.com](mailto:njasloane@gmail.com)

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.