

# Spherical Designs in Four Dimensions (Extended Abstract)

N. J. A. Sloane

AT&T Shannon Labs, Florham Park,  
NJ 07932-0971 USA  
njas@research.att.com

R. H. Hardin

227 Glenn Drive, Pataskala,  
OH 43062 USA  
rhh@cadence.com

P. Cara\*

Department of Mathematics,  
Vrije Universiteit Brussel  
Brussels, B-1050 Belgium  
pcara@vub.ac.be

February 16, 2003

**Abstract** — A preliminary report on our investigations into the existence of spherical  $t$ -designs on the unit sphere  $\Omega_4$  in 4-dimensional Euclidean space. Tables are given of the putatively best  $t$ -designs with up to 100 points. Some general constructions are proposed and explicit constructions are given for  $N$ -point strength  $t$  designs with  $N = 4p$  and  $t = \min\{p - 1, 5\}$ ,  $N = 6p$  and  $t = \min\{p - 1, 7\}$ , and  $N = 12p$  and  $t = \min\{p - 1, 11\}$ , for all  $p \geq 1$ .

## I. INTRODUCTION

Spherical  $t$ -designs have received a great deal of attention in recent years. They were the subject of a conference at the University of Geneva in October 2002, organized by Pierre de la Harpe, and there was a special session on spherical codes and designs at the annual meeting of the American Mathematical Society in Baltimore in January 2003, organized by Béla Bajnok and N.J.A.S.

Beginning in the early 1990's, the first two authors carried out a systematic search for solutions to many versions of the problem of finding the "best" way to place  $N$  points on the unit sphere  $\Omega_n$  in  $n$ -dimensional Euclidean space, concentrating on up to about 100 points in three, four and sometimes five dimensions. We originally considered just the packing problem, but the algorithm seemed to perform well on other problems, so we also tackled the covering, minimal energy and maximal volume problems, the spherical  $t$ -design problem and the problem of constructing optimal experimental designs. We also modified the algorithm to compute clusters of spheres with minimal second moment about their centroid, isometric embeddings and packings in Grassmannian spaces.

Some parts of this work have been described in [14], [15], [16], [17], [18], [28], and tables containing many of the results can be found on N.J.A.S.'s homepage [27]. The algorithm we used was a modification of the "pattern search" of Hooke and Jeeves [20]. It has been described in our earlier papers and we will say no more about it here.

Of course our computer program just produces numerical coordinates for the points  $P_1, \dots, P_N$ , typically specified to 12 decimal places. If we want to prove theorems about the results (rather than just to use them), we must show (for example) that there *is* a spherical  $t$ -design in the neighborhood of

the numerical points. We will say more about this "beautification" process later.

The purpose of the present paper is to give a preliminary report (in Section III) on the search for spherical designs in four dimensions. In Section IV we present some new constructions for four-dimensional designs that emerged from the beautification process. Full details will be published elsewhere.

## II. SPHERICAL $t$ -DESIGNS

We give five equivalent definitions.

**(D1)** A set of  $N$  points  $\{P_1, \dots, P_N\}$  on the unit sphere  $\Omega_n = S^{n-1} = \{x = (x_1, \dots, x_n) \in \mathbb{R}^n : x \cdot x = 1\}$  forms a *spherical  $t$ -design* if the identity

$$\int_{\Omega_n} f(x) d\mu(x) = \frac{1}{N} \sum_{i=1}^N f(P_i) \quad (1)$$

holds for all polynomials  $f$  of degree  $\leq t$ , where  $\mu$  is uniform measure on the sphere normalized to have total measure 1 ([10]; [12]; [9, §3.2]).

Of course a  $t$ -design is also a  $t'$ -design for all  $t' \leq t$ . The largest  $t$  for which the points form a  $t$ -design is called the *strength* of the design.

It is known that if  $N$  is large enough then a  $t$ -design in  $\Omega_n$  always exists ([25]). The problem is to find the smallest value of  $N$  for a given strength and dimension, or equivalently to find the largest strength  $t$  that can be achieved with  $N$  points in  $\Omega_n$ .

**(D2)**  $P_1, \dots, P_N$  is a spherical  $t$ -design if and only if

$$\sum_{i=1}^N f(P_i) = 0 \quad (2)$$

for all harmonic polynomials  $f$  of degrees from 1 to  $t$ . ( $f$  is harmonic if the Laplacian

$$\Delta f = \sum_{j=1}^n \frac{\partial^2 f}{\partial x_j^2}$$

vanishes.)

\*Postdoctoral fellow of the Fund for Scientific Research-Flanders (Belgium). Philippe Cara thanks AT&T Shannon Labs for support during his visit in Fall 2002

(D3)  $P_1, \dots, P_N$  is a spherical  $t$ -design if and only if

$$\sum_{i=1}^N f(P_i^T) = \sum_{i=1}^N f(P_i) \quad (3)$$

for all polynomials  $f$  of degrees from 1 to  $t$  and all orthogonal transformations  $T \in O(n, \mathbb{R})$ .

(D4) For  $\alpha \in [-1, 1]$ , let  $A_\alpha$  be the number of ordered pairs  $(i, j)$  such that  $P_i \cdot P_j = \alpha$ . The numbers  $\{A_\alpha\}$  are the *distance distribution* of the design. Then the Gegenbauer transform

$$\sum_{\alpha} A_{\alpha} Q_k(\alpha) \geq 0 \text{ for all } k, \quad (4)$$

and  $P_1, \dots, P_N$  is a spherical  $t$ -design if and only if equality holds in (4) for all  $k = 1, \dots, t$ . The  $Q_k(x)$  are Gegenbauer or ultraspherical polynomials defined by the recurrence

$$Q_0 = 1, \quad Q_1 = nx,$$

$$\lambda_{k+1} Q_{k+1} = x Q_k - (1 - \lambda_{k-1}) Q_{k-1},$$

where  $\lambda_k = k/(n + 2k - 2)$ .

The inequalities (4) are the basis for the linear programming bound for spherical designs.

(D5)  $P_1, \dots, P_N$  forms a spherical  $t$ -design if and only if the polynomial identities

$$\frac{1}{N} \sum_{i=1}^N (P_i \cdot x)^{2s} = \left( \prod_{j=0}^{s-1} \frac{2j+1}{2j+n} \right) (x \cdot x)^s, \quad (5)$$

and

$$\frac{1}{N} \sum_{i=1}^N (P_i \cdot x)^{2\bar{s}+1} = 0,$$

hold, where  $s$  and  $\bar{s}$  are defined by  $\{2s, 2\bar{s} + 1\} = \{t - 1, t\}$  ([12]; [23, p. 114], [24]).

**Spherical  $t$ -designs from groups.** Let  $G$  be a finite subgroup of  $O(n, \mathbb{R})$ . Consider the ring  $R$  of polynomials in  $x_1, \dots, x_n$  that are invariant under  $G$ , ignoring the trivial invariant  $x_1^2 + \dots + x_n^2$ . If  $R_j$  is the linear subspace of  $R$  consisting of homogeneous invariants of degree  $j$ , then the dimensions  $d_j = \dim R_j$  are given by the Molien series

$$\Phi_G(\lambda) = \frac{1}{|G|} \sum_{T \in G} \frac{1 - \lambda^2}{\det(I - \lambda T)} = \sum_{j=0}^{\infty} d_j \lambda^j. \quad (6)$$

Let  $\mathcal{P} = \{P_1, \dots, P_N\}$  be a union of  $m$  orbits under  $G$  (not necessarily all of the same size). There are  $m(n - 1)$  degrees of freedom in choosing  $\mathcal{P}$ .

Then  $\mathcal{P}$  is a spherical  $t$ -design if and only if the average of  $f$  over  $\mathcal{P}$  is equal to the average of  $f$  over  $\Omega_n$  for all  $f \in R_1 \cup R_2 \cup \dots \cup R_t$  [11], [13], [19]. This imposes a total of  $e_t = d_1 + \dots + d_t$  conditions on  $\mathcal{P}$ . So if  $m(n - 1) \geq e_t$ , we may hope that we can choose the orbits to form a  $t$ -design.

In particular, suppose that  $d_1 = \dots = d_t = 0$  for some  $t$ . Then there are no conditions to be satisfied, and so every orbit

under  $G$  is a  $t$ -design. (This result seems to be due to Sobolev [29], cf. [9].)

Furthermore, if  $d_{t+1} = 1$ , and  $\psi$  is the unique (up to scalars) invariant of degree  $m$ , then the orbit of any real zero of  $\psi$  is at least a  $(t + 1)$ -design.

### III. FOUR DIMENSIONS

Table 1 gives our lower bounds for  $\tau_4(N)$ . The third column gives the order of the largest symmetry group we have found for a design with the specified strength. **Bold** entries indicate where  $\tau_4(N)$  appears to exceed  $\tau_4(k)$  for all  $k < N$ .

**Notes on Table 1.** (i) For each entry in the table we have a numerical approximation to the claimed design; that is, numerical coordinates for points such that the magnitude of the difference between the left and right sides of (5) is less than  $10^{-8}$ . These designs will be placed on N.J.A.S.'s home page [27].

(ii) The entries with  $N \leq 20, 24, 48$  and  $120$  were previously known to exist (Mimura [21], Bajnok [2], [3], Hardin and Sloane [14], Delsarte, Goethals and Seidel [10]); and the entries with  $N = 21 - 23$  and  $25, 26, 27, 31$  were previously conjectured to exist [14].

(iii) In the present investigation the entries with  $N = 21, 26, 28, 32, 36, 39, 40, 42, 44, 52, 54, 60, 65, 66, 72, 78, 84, 90$  have been beautified (and so shown to exist).

(iv) Many entries still have not yet been beautified, for example those with  $N = 22, 23, \dots, 30, \dots$  points, and especially the putative 8-design with  $N = 96$  points. (The biggest group we have found for such a design has order only 6, with 16 orbits, which leaves too many unknown parameters for our present beautification methods to handle.)

(v) For definitions of orthoplex, diplo-simplex, etc., see [8].

(vi) This is only a preliminary version of the table. It is possible that some entries — especially the group orders — may change in the final version.

The data in Table 1 can be summarized as follows. It appears that spherical designs in four dimensions with strength  $t$  and  $N$  points exist if and only if:

$t$	$N$
2	$\geq 5$
3	$8, \geq 10$
4	$\geq 20$
5	$24, 28, 30, \geq 32$
6	$42, 48, \geq 50$
7	$48, 54, 56, 60, 62, 64, 66, \geq 68$
8	$\geq 96$
...	...

For  $t = 2$  this is known to be true from the work of Mimura [21], and for  $t = 3$  from Bajnok [2], [3] and Boyvalenkov, Danev and Nikova [4]. For  $t \geq 4$  these are only conjectures.

**Examples.** Our beautification technique begins by computing the distance distribution of the points, their apparent automorphism group and its orbits on the points.

For small numbers of points this is often enough to reveal the structure immediately.

**7 points, strength 2.** The group has order 14 and acts transitively on the points. By a suitable change of coordinates we can make the generators for the group look like

$$\begin{bmatrix} \cos \theta_1 & \sin \theta_1 & 0 & 0 \\ -\sin \theta_1 & \cos \theta_1 & 0 & 0 \\ 0 & 0 & \cos \theta_2 & \sin \theta_2 \\ 0 & 0 & -\sin \theta_2 & \cos \theta_2 \end{bmatrix} \quad (7)$$

and  $\text{diag}\{1, -1, 1, -1\}$ , where  $\theta_1 = 2\pi/7$ ,  $\theta_2 = 4\pi/7$ . This transformation also beautifies the points, which become

$$\frac{1}{\sqrt{2}}(\cos j\theta, \sin j\theta, \cos 2j\theta, \sin 2j\theta)$$

( $j = 0, \dots, 6$ ). We may specify this design more simply as  $\{(\omega^j, \omega^{2j}), j = 0, \dots, 6\}$ , where  $\omega = e^{2\pi i/7}$ , which is how it is listed in Table 1. Once we have these simple coordinates it is straightforward to use a symbolic algebra package such as Maple [6] to verify that the strength is as claimed.

Of course this design has been known for a long time, but we have described the process in detail to illustrate the beautification technique.

The designs for  $N = 9$  and 13 are of the same type.

**20 points, strength 4.** This was already beautified in [14]. The points consist of the orbit of a single point  $(a, b, c, d)$  under a certain group of order 20. The equations that  $a, b, c, d$  must satisfy are given in [14]. At that time we found that  $a, b, c, d$  were roots of integral polynomials of degree 72. We have now found that these polynomials can be factored, and so a simpler description is possible. It turns out that  $c = -0.4594335\dots$  is a root of the following polynomial of degree 20 over  $\mathbf{Q}[\sqrt{3}]$ :

$$\begin{aligned} &1019215872 x^{20} - 2548039680 x^{18} \\ &+ 849346560 x^{18}\sqrt{3} + 3609722880 x^{16} \\ &- 1804861440 x^{16}\sqrt{3} - 3185049600 x^{14} \\ &+ 1769472000 x^{14}\sqrt{3} + 1761177600 x^{12} \\ &- 1006387200 x^{12}\sqrt{3} - 608587776 x^{10} \\ &+ 350631936 x^{10}\sqrt{3} + 128240640 x^8 \\ &- 74108160 x^8\sqrt{3} - 15465600 x^6 \\ &+ 8950080 x^6\sqrt{3} + 930300 x^4 \\ &- 538800 x^4\sqrt{3} - 20100 x^2 \\ &+ 11650 x^2\sqrt{3} - 63 + 36\sqrt{3}, \end{aligned}$$

and  $a, b$  and  $d$  are roots of similar polynomials of the same degree.

A different 20-point 4-design will be given in Section V.

**21 points, strength 4.** The group of this design is the same as the group of the above 7-point design. The 21 points fall into two orbits, with representatives  $(u, v, w, x)$  and  $(y, 0, z, 0)$  (say), whose approximate values are

$$(0.01210\dots, -0.83898\dots, -0.04657\dots, -0.54202\dots)$$

and

$$(0.30292\dots, 0, -0.95301\dots, 0).$$

After a considerable amount of simplification we find that the following equations must be satisfied for this to be a 4-design:

$$\begin{aligned} u^2 + v^2 + w^2 + x^2 &= y^2 + z^2 = 1, \\ u^2 + v^2 &= \frac{1}{4}(3 - 2y^2), \\ (u^2 + v^2)^2 &= \frac{1}{2}(1 - y^4), \\ (u^2 + v^2)(w^2 + x^2) &= \frac{1}{4}(1 - 2y^2z^2), \\ (w^2 + x^2)^2 &= \frac{1}{2}(1 - z^4), \\ 2w(u^2 - v^2) + 4uvx &= -y^2z, \\ 6wx(ux + vw) - 2uw^3 - 2vx^3 &= -yz^3. \end{aligned}$$

These imply

$$12y^4 - 12y^2 + 1 = 0,$$

$$u^2 + v^2 = \frac{1}{2} + \frac{1}{2\sqrt{6}},$$

and finally that  $u$  is a root of the irreducible polynomial

$$\begin{aligned} &370064301985047250503374330658816000000 u^{56} \\ &- 2590450113895330753523620314611712000000 u^{54} \\ &+ 8499914436219054034999379157319680000000 u^{52} \\ &- 17391094775057715319228888466325504000000 u^{50} \\ &+ 24899387662279537368324714887380992000000 u^{48} \\ &- 26538129846554061588093258386374656000000 u^{46} \\ &+ 21869812340899414041740569007357952000000 u^{44} \\ &- 14288148063124617412183158428054467903488 u^{42} \\ &+ 7528498207750245383367572061822929338368 u^{40} \\ &- 3237480950055173571219844766822842761216 u^{38} \\ &+ 1145499747167940743157516608475821506560 u^{36} \\ &- 335215964942931989772955061161455255552 u^{34} \\ &+ 81357491663702586946618517659406303232 u^{32} \\ &- 16386240716737992044502246170623475712 u^{30} \\ &+ 2734899456377829726261221866315186176 u^{28} \\ &- 376908341370344979393267774145953792 u^{26} \\ &+ 42638421653218179958830238871322624 u^{24} \\ &- 3925374570079154331047708023848960 u^{22} \\ &+ 290546738656133501034283042603008 u^{20} \\ &- 17003134241759862086722092269568 u^{18} \\ &+ 768504591189193729587321765888 u^{16} \\ &- 25936049223509962319342862336 u^{14} \\ &+ 620859209067240422756843520 u^{12} \\ &- 9680230988740463748710400 u^{10} \\ &+ 83470588537572204192000 u^8 \\ &- 264480246652176979200 u^6 \\ &+ 139741070071159200 u^4 \\ &- 24494390617200 u^2 \\ &+ 1383468025. \end{aligned}$$

#### IV. A NEW CONSTRUCTION

The most interesting example that we beautified was the 6-design in  $\Omega_4$  with 42 points. This turned out to consist of six heptagons, each in a different plane, with a group of order 42 acting transitively on the 42 points. Or, to put it another way: the computer-produced design was suggesting to us that we

should choose six planes in  $\mathbb{R}^4$ , i.e. six points in the Grassmann manifold  $G(4, 2)$ , and draw a heptagon in each plane.

This was an appealing idea, in view of the recent work on finding packings and designs in Grassmann manifolds (cf. [1], [5], [7], [26])! It immediately suggested a general construction.

Informally stated, we take a “nice” set of  $M$  planes in  $\mathbb{R}^4$ , draw a regular  $p$ -gon in each plane for a given value of  $p$ , obtaining a set of  $N = Mp$  points that will form a spherical  $t$ -design for some  $t$  (possibly 0).

The following are two more precise versions.

**Construction 1.** Fix integers  $M$  and  $p$ , and let  $\theta = 2\pi/p$ . Let  $\Pi_i$  ( $i = 1, \dots, M$ ) be a plane in  $\mathbb{R}^4$  spanned by a pair of orthogonal vectors  $u_i, v_i$ , and let  $\phi_1, \dots, \phi_M \in [0, 2\pi)$  be phase angles. Then the design consists of the  $N = Mp$  points

$$\{\cos(j\theta + \phi_i)u_i + \sin(j\theta + \phi_i)v_i : 0 \leq j < p, 1 \leq i \leq M\}. \quad (8)$$

**Construction 2.** Fix integers  $M$  and  $p$ , and let  $\theta = 2\pi/p$ . For  $i = 1, \dots, M$  let  $\Pi_i = \langle u_i, v_i \rangle$ ,  $\Pi'_i = \langle w_i, x_i \rangle$  be a pair of perpendicular planes in  $\mathbb{R}^4$ , where  $\{u_i, v_i, w_i, x_i\}$  is a coordinate frame, let  $\phi_1, \dots, \phi_M \in [0, 2\pi)$  be phase angles, and let  $k_1, \dots, k_M$  be integers. Then the design consists of the  $N = Mp$  points

$$\begin{aligned} &\{\cos(j\theta + \phi_i)u_i + \sin(j\theta + \phi_i)v_i \\ &\quad + \cos(jk_i\theta + \phi_i)w_i + \sin(jk_i\theta + \phi_i)x_i : \\ &\quad 0 \leq j < p, 1 \leq i \leq M\}. \end{aligned} \quad (9)$$

The first construction generalizes the 42-point design and the second construction generalizes the 7-, 9- and 13-point designs (and others).

We are far from completing our investigations of these constructions. At present three concrete examples have emerged, applications of Construction 1 with all phase angles  $\phi_i$  set to 0. (There are also simpler applications of Construction 1 using a pair of planes to obtain 3-designs, which we will not describe here.)

**Theorem 1** Let  $\Pi_1, \dots, \Pi_4$  be the planes with generator matrices

$$\begin{aligned} &\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & s & s & s \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 & 0 \\ s & 0 & s & -s \end{bmatrix}, \\ &\begin{bmatrix} 0 & 0 & 1 & 0 \\ s & -s & 0 & s \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 & 1 \\ s & s & -s & 0 \end{bmatrix}, \end{aligned}$$

where  $s = 1/\sqrt{3}$ . Then (8) (with all  $\phi_i = 0$ ) forms a  $4p$ -point spherical  $t$ -design with  $t = \min\{p - 1, 5\}$ .

Theorem 1 yields the following  $t$ -designs:

$p$	2	3	4	5	6	7	8	9	10	11	12	...
$N$	8	12	16	20	24	28	32	36	40	44	48	...
$t$	3	2	3	4	5	5	5	5	5	5	5	...
group	384	18	96	15	1152	21	192	27	240	33	288	...

**Notes.** (i)  $p = 2$  gives the orthoplex, which has higher strength than was guaranteed by the theorem. (ii)  $p = 5$  gives a simpler construction for a 20-point 4-design than the one in [14]. (iii)  $p = 6$  gives the 24-cell. (iv) The designs are antipodal if and only if  $p$  is even, and this is reflected in the fact that the designs with  $p$  odd have much smaller groups.

**Theorem 2** Let  $\Pi_1, \dots, \Pi_6$  be the planes with generator matrices

$$\begin{aligned} &\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \\ &\begin{bmatrix} s & 0 & -h & h \\ 0 & s & -h & -h \end{bmatrix}, \begin{bmatrix} -h & h & s & 0 \\ -h & -h & 0 & s \end{bmatrix}, \\ &\begin{bmatrix} -s & 0 & -h & -h \\ 0 & -s & h & -h \end{bmatrix}, \begin{bmatrix} -h & -h & -s & 0 \\ h & -h & 0 & -s \end{bmatrix}, \end{aligned}$$

where now  $s = 1/\sqrt{2}$ ,  $h = 1/2$ . Then (8) (with all  $\phi_i = 0$ ) forms a  $6p$ -point  $t$ -design with  $t = \min\{p - 1, 7\}$ .

Theorem 2 yields the following  $t$ -designs:

$p$	2	3	4	5	6	7	8	9	10	11	12	...
$N$	12	18	24	30	36	42	48	54	60	66	72	...
$t$	1	2	3	4	5	6	7	7	7	7	7	...
group	24	36*	128*	30	144*	42	2304	54	120	66	192*	...

**Notes.** (i) The starred group orders were obtained using a slightly different set of six planes. (ii)  $p = 6$  is the 42-point design that the computer found. (iii)  $p = 8$  gives the 48-point design formed from two copies of the 24-cell. Its group is the Clifford group  $\mathcal{C}(2)$  from [22].

We remind the reader that there is a canonical way to specify a plane  $\Pi$  in  $\mathbb{R}^4$  by a pair of points  $(l, r) \in \Omega_3 \times \Omega_3$  (see [7], Theorem 1). A set of  $M$  planes is described by a “binocular spherical code”  $\{(l_i, r_i) \in \Omega_3 \times \Omega_3 : i = 1, \dots, M\}$ . The sets of planes used in Theorems 1 and 2 correspond to the binocular spherical codes {point  $\times$  tetrahedron} and {point  $\times$  octahedron}. The third theorem is a natural continuation:

**Theorem 3** By using the  $M = 12$  planes described by {point  $\times$  icosahedron} we obtain from (8) a  $12p$ -point spherical  $t$ -design with  $t = \min\{p - 1, 11\}$ .

Theorem 3 yields the following  $t$ -designs:

$p$	6	7	8	9	10	11	12	13	14	15	...
$N$	72	84	96	108	120	132	144	156	168	180	...
$t$	5	6	7	8	9	10	11	11	11	11	...
group	24	28	32	36	40	44	48	52	56	60	...

Theorems 1–3 are established by computing the distance distribution of the design and working out its Gegenbauer transform (cf. (4)). This is simplified by the fact that the planes in the three theorems form isoclinic sets (cf. [7]). Details are postponed to the full version of this paper.

## REFERENCES

[1] C. Bachoc, R. Coulangéon and G. Nebe, Designs in Grassmannian spaces and lattices, *J. Algebraic Combin.*, **16** (2002), 5–19.

- [2] B. Bajnok, Construction of spherical  $t$ -designs, *Geom. Dedicata*, **43** (1992), 167–179.
- [3] B. Bajnok, Construction of spherical 3-designs, *Graphs and Combinatorics*, **14** (1998), 97–107.
- [4] P. Boyvalenkov, D. Danev and S. Nikova, Nonexistence of certain spherical designs of odd strengths and cardinalities, *Discrete Comput. Geom.*, **21** (1999), 143–156.
- [5] A. R. Calderbank, R. H. Hardin, E. M. Rains, P. W. Shor and N. J. A. Sloane, A Group-Theoretic Framework for the Construction of Packings in Grassmannian Spaces, *J. Algebraic Combin.*, **9** (1999), 129–140.
- [6] B. W. Char et al., *Maple V Reference Manual*, Springer-Verlag, NY, 1991.
- [7] J. H. Conway, R. H. Hardin, and N. J. A. Sloane, Packing lines, planes, etc.: packings in Grassmannian space, *Experimental Math.*, **5** (1996), 139–159.
- [8] J. H. Conway and N. J. A. Sloane, The cell structures of lattices, in *Miscellanea mathematica*, P. Hilton et al., editors, Springer-Verlag, New York, 1991, pp. 71–107.
- [9] J. H. Conway and N. J. A. Sloane, *Sphere Packings, Lattices and Groups*, Springer-Verlag, New York, 3rd edition, 1999.
- [10] P. Delsarte, J.-M. Goethals and J. J. Seidel, Spherical codes and designs, *Geom. Dedicata*, **6** (1977), 363–388.
- [11] J.-M. Goethals and J. J. Seidel, Spherical designs, in D. K. Ray-Chaudhuri, ed., *Relations Between Combinatorics and Other Parts of Mathematics, Proc. Symp. Pure Math.*, **34** (1979), 255–272.
- [12] J.-M. Goethals and J. J. Seidel, Cubature formulae, polytopes and spherical designs, in C. Davis et al., eds., *The Geometric Vein: The Coxeter Festschrift*, Springer-Verlag, NY, 1981, pp. 203–218.
- [13] J.-M. Goethals and J. J. Seidel, The football, *Nieuw Archief Wisk.*, **29** (1981), 50–58.
- [14] R. H. Hardin and N. J. A. Sloane, New spherical 4-designs, *Discrete Mathematics*, **106/107** (1992), 255–264. (Topics in Discrete Mathematics, vol. 7, "A Collection of Contributions in Honour of Jack Van Lint", ed. P. J. Cameron and H. C. A. van Tilborg, North-Holland, 1992.)
- [15] R. H. Hardin and N. J. A. Sloane, A new approach to the construction of optimal designs, *J. Statistical Planning and Inference*, Vol. 37, 1993, pp. 339–369.
- [16] R. H. Hardin and N. J. A. Sloane, *Operating Manual for Gosset: A General-Purpose Program for Constructing Experimental Designs (Second Edition)*, Statistics Research Report No. 98, AT&T Bell Labs, Murray Hill, NJ, Nov. 15, 1991. Also DIMACS Technical Report 93–51, August 1993, Center for Discrete Math. and Computer Science, Rutgers Univ., New Brunswick, NJ.
- [17] R. H. Hardin and N. J. A. Sloane, Expressing  $(a^2 + b^2 + c^2 + d^2)^2$  as a sum of 23 sixth powers, *J. Combin. Theory, Series A*, **68** (1994), 481–485.
- [18] R. H. Hardin and N. J. A. Sloane, Codes (Spherical) and Designs (Experimental), in *Different Aspects of Coding Theory*, edited A. R. Calderbank, AMS Series Proceedings Symposia Applied Math., Vol. **50**, 1995, pp. 179–206.
- [19] R. H. Hardin and N. J. A. Sloane, McLaren's Improved Snub Cube and Other New Spherical Designs in Three Dimensions, *Discrete Comput. Geometry*, **15** (1996), 429–441.
- [20] R. Hooke and T. A. Jeeves, 'Direct search' solution of numerical and statistical problems, *J. Assoc. Computing Machinery*, **8** (1961), 212–229.
- [21] Y. Mimura, A construction of spherical 2-designs, *Graphs and Combinatorics*, **6** (1990), 369–372.
- [22] G. Nebe, E. M. Rains and N. J. A. Sloane, The Invariants of the Clifford Groups, *Designs, Codes and Cryptography* **24** (2001),
- [23] B. Reznick, *Sums of Even Powers of Real Linear Forms*, Memoirs Amer. Math. Soc., No. 463, March 1992.
- [24] B. Reznick, Some constructions of spherical 5-designs, *Linear Algebra and Its Applications*, **226/228** (1995), 163–196.
- [25] P. D. Seymour and T. Zaslavsky, Averaging sets: a generalization of mean values and spherical designs, *Advances in Math.*, **52** (1984), 213–240.
- [26] P. W. Shor and N. J. A. Sloane, A family of optimal packings in Grassmannian manifolds, *J. Algebraic Combin.*, **7** (1998), 157–163.
- [27] N. J. A. Sloane, Home page, <http://www.research.att.com/~njas/>, 2003.
- [28] N. J. A. Sloane, R. H. Hardin, T. S. Duff and J. H. Conway, Minimal-energy clusters of hard spheres, *Discrete Comput. Geom.*, **14** 1995, 237–259.
- [29] S. L. Sobolev, Cubature formulae on the sphere invariant under finite groups of rotations (Russian), *Doklady Akademii Nauk SSR*, **146** (No. 2, 1962), 310–313; translation in *Soviet Mathematics Doklady*, **3** (1962), 1307–1310.

Table 1: Conjectured values of  $\tau_4(N)$ , the largest  $t$  for which an  $N$ -point configuration on the sphere in 4 dimensions forms a spherical  $t$ -design.

$N$	$\tau_4(N)$	Group	Remarks	$N$	$\tau_4(N)$	Group	Remarks
1	<b>0</b>	$\infty$	single point	52	6	52	
2	<b>1</b>	$\infty$	two antipodal points	53	6	6	
3	1	$\infty$	equilateral triangle	54	7	54	Theorem 2
4	1	48	tetrahedron	55	6	20	
5	<b>2</b>	120	regular simplex	56	7	16	
6	2	72	join of 2 triangles	57	6	9	
7	2	14	$\{(\omega^j, \omega^{2j})\}$	58	6	116	
8	<b>3</b>	384	orthoplex	59	6	12	
9	2	72	$\{(\omega^j, \omega^k)\}$	60	7	240	Theorem 2
10	3	240	diplo-simplex	61	6	8	
11	3	22		62	7	8	
12	3	288	join of 2 hexagons	63	6	18	
13	3	52	$\{(\omega^j, \omega^{5j})\}$	64	7	256	
14	3	392		65	6	52	
15	3	60		66	7	66	Theorem 2
16	3	512	join of 2 octagons	67	6	12	
17	3	68		68	7	136	
18	3	648		69	7	3	
19	3	48		70	7	40	
20	<b>4</b>	20	Theorem 1; [14]	71	7	6	
21	4	14		72	7	576	Theorem 2
22	4	6		73	7	6	
23	4	6		74	7	16	
24	<b>5</b>	1152	24-cell	75	7	6	
25	4	20		76	7	38	
26	4	52		77	7	22	
27	4	18		78	7	52	
28	5	28	Theorem 1	79	7	8	
29	4	8		80	7	256	
30	5	40		81	7	36	
31	4	12		82	7	164	
32	5	192	Theorem 1	83	7	12	
33	5	6		84	7	144	
34	5	12		85	7	34	
35	5	10		86	7	24	
36	5	144	Theorem 1	87	7	18	
37	5	12		88	7	128	
38	5	48		89	7	12	
39	5	26		90	7	180	
40	5	240	Theorem 1	91	7	52	
41	5	12		92	7	48	
42	<b>6</b>	42	Theorem 2	93	7	36	
43	5	12		94	7	96	
44	5	96	Theorem 1	95	7	38	
45	5	30		96	<b>8</b>	6	
46	5	96		97	8	1	
47	5	12		98	8	4	
48	<b>7</b>	2304	two 24-cells	99	8	11	
49	5	4		100	8	20	
50	6	100		...	...		
51	6	4		120	<b>11</b>	14400	600-cell [10]